

A Fallback Disk Accretion-Involved Formation Channel to PSR J1903+0327

Xi-Wei Liu¹ and Xiang-Dong Li²

liuxw@phy.ccnu.edu.cn; lixd@nju.edu.cn

ABSTRACT

The discovery of the eccentric binary and millisecond pulsar PSR J1903+03273 has raised interesting questions about the formation mechanisms of this peculiar system. Here we present a born-fast scenario for PSR J1903+03273. We assume that during the supernova (SN) explosion that produced the pulsar, a fallback disk was formed around and accreted onto the newborn neutron star. Mass accretion could accelerate the neutron star's spin to milliseconds, and decrease its magnetic field to $\sim 10^8 - 10^9$ G, provided that there was sufficient mass ($\sim 0.1M_\odot$) in the fallback disk. The neutron star became a millisecond pulsar after mass accretion terminated. In the meanwhile the binary orbit has kept to be eccentric (due to the SN explosion) for $\sim 10^9$ yr. We have performed population synthesis calculations of the evolutions of neutron stars with a fallback disk, and found that there might be tens to hundreds of PSR J1903+03273-like systems in the Galaxy. This scenario also suggests that some fraction of isolated millisecond pulsars in the Galactic disk could be formed through the same channel.

Subject headings: accretion, accretion disks — pulsars: general — pulsars: individual (PSR J1903+0327) — stars: neutron

1. Introduction

There are now around 100 binary and millisecond pulsars (BMPSRs) known in the Galaxy. Most of them are characterized by short spin periods ($P \lesssim 15$ ms), weak magnetic fields ($B \sim 10^8 - 10^9$ G), and circular orbits with companions of mass $\sim 0.15M_\odot - 0.45M_\odot$. In the recycling scenario, these systems are thought to be the descendants of low-mass X-ray binaries (Bhattacharya & van den Heuvel 1991; Tauris & van den Heuvel 2006). The

¹Institute of Astrophysics, Huazhong Normal University, Wuhan 430079, China

²Department of Astronomy, Nanjing University, Nanjing 210093, China

evolutionary path is briefly described as follows. A high-field ($B \sim 10^{12} - 10^{13}$ G), rapidly rotating neutron star (NS) is born in a binary with a low-mass ($\sim 1M_{\odot}$) main-sequence (MS) companion star. During the supernova (SN) that produces the NS, mass loss and a kick imparted on the NS cause the orbit to be eccentric. The NS spins down as a radio pulsar for $\sim 10^6 - 10^7$ yr until passing by the so-called “death line” in the magnetic field - spin period ($B - P$) diagram (Ruderman & Sutherland 1975). When the companion evolves to overflow its Roche lobe, mass transfer occurs by way of an accretion disk, and tidal friction serves to circularize the orbit. Mass accretion onto the NS gives rise to X-ray emission, induces magnetic field decay (the mechanisms for the field decay induced by accretion are, however, not well understood), and accelerates the NS up to short (millisecond) period. When the companion loses almost all of its envelope and mass transfer ceases, the endpoint of the evolution is a circular binary containing an NS visible as a low-field, millisecond radio pulsar, and a He or CO white dwarf (WD), the remaining core of the companion. A fast-growing population of eccentric ($e > 0.1$) BMPSRs have been recently revealed in globular clusters (GCs) (Freire, Ransom & Gupta 2007). These high eccentricities are likely to be attributed to the dynamical interactions in the central regions of GCs.

Most recently Champion et al. (2008) reported the discovery of an eccentric BMPSR PSR J1903+0327 in the Galactic plane. With a spin period of 2.15 ms, the pulsar lies in an eccentric ($e = 0.44$), 95-day orbit around a $\sim 1M_{\odot}$ companion. The mass of the pulsar was estimated to be $1.74 \pm 0.04 M_{\odot}$, about 30% larger than that of other binary NSs in the Galactic disk. Infrared observations identified a possible MS companion star. Preferred formation scenarios may include (cf. van den Heuvel 2008; Champion et al. 2008): (1) the pulsar was recycled in a GC, then the original donor star was replaced by the present MS companion via one or more exchange interactions, and the binary was displaced from the GC due to either ejection or the disruption of the GC; (2) the pulsar is part of a primordial hierarchical triple system, recycled by accretion from the progenitor of a massive ($\sim 0.9 - 1.1 M_{\odot}$) WD which is seen in the timing measurement, while the detected infrared counterpart, the third star, is in a much wider and highly inclined orbit around the inner binary. The eccentricity of the inner binary is caused by the perturbation of the outer MS star; (3) in an alternative triple-star model, the WD in the inner binary system was evaporated by the pulsar’s energy flux, or coalesced with it, such that the present binary remained.

Since there is no strong evidence for the GC or triple star origin of PSR J1903+0327, in this paper, we seek an alternative, born-fast scenario for PSR J1903+0327 which was already mentioned by Champion et al. (2008). These authors argued that the pulsar was not likely to be formed spinning rapidly at the time of SN with a small magnetic field, due to the following reasons: (1) there are no pulsars like PSR J1903+0327 in any of the > 50 young

supernova remnants in which an NS has been inferred or detected directly; (2) the 18 isolated MPSRs detected in the Galactic disk have spin distributions, space velocities, and energetics indistinguishable from those of recycled BMPRSs and their space velocities and scale heights do not match those of non-recycled pulsars; (3) magnetic fields in young pulsars is unlikely to be less than 10^{10} G. However, as we shown below, if experienced accretion from a SN fallback disk, the newborn pulsar could be accelerated to a period of several milliseconds, together with the magnetic field decayed to $\sim 10^8$ G from an initial value, say $\sim 10^{12}$ G. When the disk becomes neutral after $\sim 10^3 - 10^4$ yr (Menou, Perna & Hernquist 2001) and accretion ceases, the NS becomes an MPSR. The orbit, if not being disrupted by the SN explosion, will remain eccentric for $> 10^{10}$ yr before it is circularized (cf. Hurley, Tout & Pols 2002). The binary consists of a recycled pulsar and a MS companion in an eccentric orbit, just like PSR J1903+0327. Hence this scenario could produce a population of eccentric BMPRSs as well as isolated MPSRs.

2. Spin Evolution of An NS with A Fallback Disk

Following the formation of an NS through the core collapse of its progenitor, a small amount of mass could fall back onto the compact object (Colgate 1971; Chevalier 1989; Lin, Woosley & Bodenheimer 1991). Some of the fallback material may carry sufficient angular momentum and to form an accretion disk of mass M_d around the NS. The initial transient phase lasts for a local viscous timescale $t_0 \approx 6.6 \times 10^{-5} (T_{c,6})^{-1} R_{d,8}^{1/2} (t_0)$ yr, on which a thin disk forms (Cannizzo, Lee & Goodman 1990; Menou, Perna & Hernquist 2001). Here $T_{c,6}$ is the typical temperature in the outermost disk annulus during this early phase, in units of 10^6 K, and $R_{d,8}(t_0)$ is the initial disk radius in units of 10^8 cm. We set $T_{c,6} = 1$ throughout this paper. The subsequent evolution of the disk obeys the self-similar solution (Cannizzo, Lee & Goodman 1990), i.e.,

$$\dot{M}_d(t) = \begin{cases} \dot{M}_d(t_0), & 0 < t < t_0, \\ \dot{M}_d(t_0) \left(\frac{t}{t_0}\right)^{-p}, & t \geq t_0, \end{cases} \quad (1)$$

where the power index $p = 1.25$ (Francischelli & Wijers 2002), $\dot{M}_d(t_0)$ is a constant, which we normalize to the total mass of the disk $M_d = \int_0^\infty \dot{M}_d dt$, by $M_d/(5t_0)$ (Chatterjee, Hernquist & Narayan 2000).

The subsequent evolution of the NS can be divided into three phases:

1. The *accretor* phase - When the magnetosphere radius $R_m \simeq 1.6 \times 10^8 B_{12}^{4/7} \dot{M}_{18}^{-2/7}$ cm is less than the corotation radius defined by $R_c = (GM/\Omega)^{1/3}$, the fallback matter is allowed to be accreted onto the surface of the NS. Here $B_{12} = B/10^{12}$ G, $\dot{M}_{18} = \dot{M}/10^{18}$ gs $^{-1}$, and

Ω is the angular velocity of the NS. To evaluate the accretion torque, we divide this phase further into two sub-phases: a) If $R_m \leq$ the NS radius R_{NS} , the torque exerted on the star is assumed to be $\dot{J} = \dot{M} R_{NS}^2 \Omega_K(R_{NS})$, where $\Omega_K(R)$ is the Keplerian velocity at R ; b) if $R_{NS} < R_m \leq R_c$, the accretion torque is given by $\dot{J} = 2\dot{M} R_m^2 \Omega_K(R_m)[1 - \Omega/\Omega_K(R_m)]$.

The accretion rate $\dot{M}$ of the NS is assumed to be limited by the Eddington accretion rate $\dot{M}_E \simeq 1 \times 10^{18} \text{ g s}^{-1}$ for a $1.4 M_\odot$ NS. However, during the early phase of accretion, the accretion rate may be so high that the radiation is trapped in the flow and neutrino losses are important close to the NS surface (Colgate 1971; Chevalier 1989). The gravitational energy generated by infalling matter is mainly released by neutrino losses. Hence the Eddington limit does not work in this phase. According to Chevalier (1989), when the accretion rate drops to about $3 \times 10^{-4} M_\odot \text{ yr}^{-1}$, the reverse shock reaches the radiation trapping radius so that the photons can begin to diffuse out from the shocked envelope. A luminosity of Eddington limit is expected now, and this luminosity can reduce or even reverse the inflow outside the shocked envelope. Hence we adopt the Eddington limited accretion only when $\dot{M} < 3 \times 10^{-4} M_\odot \text{ yr}^{-1}$. Otherwise we assume all the mass is accreted by the NS.

Along with mass accretion, we assume that the magnetic field evolution follows the relation, $B = B_0(1 + \Delta M/M_0)$, where B_0 is the initial magnetic field strength, and M_0 is a parameter that determines the rate of decay with its typical value of $\sim 10^{-5} - 10^{-4} M_\odot$ (Taam & van den Heuvel 1986; Shibazaki et al. 1989; Romani 1990).

2. The *propeller* phase - This phase begins when $\dot{M}$ decreases so that $R_c < R_m$. The infalling matter is assumed to be accelerated outward owing to the centrifugal barrier, taking away the angular momentum of the NS (Illarinov & Sunyaev 1975). In this phase we use the same formula as in the *accretor* phase b) to estimate the propeller spin-down torque.

3. The *radio pulsar* phase - As $\dot{M}$ decreases further, R_m exceeds the light cylinder radius $R_{lc} = c/\Omega$, or the disk becomes neutral, the accretion ceases and the NS becomes a radio pulsar. The neutral timescale of a fallback disc is $t_n = [R_{d,10}^3(t_0) \times 10^{16} \text{ g s}^{-1} / \dot{M}_d(t_0)]^{-1/2.75} t_0$ (Menou, Perna & Hernquist 2001), where $R_{d,10}$ is the disk radius in units of 10^{10} cm . In this work, when either $R_m > R_{lc}$ or $t > t_n$, we assume that the radio pulsar phase turns on and the subsequent NS spin evolution follows the magnetic dipole radiation prescription, until the NS crosses the "death line", i.e. $B_{12}/P^2 < 0.17$.

In Fig. 1 we show several examples of the calculated evolutions of the NS spin and magnetic field under typical conditions according the scheme described above. The model parameters are listed in Table 1. For the spin evolution, the thick solid, solid, and dotted lines represent the *accretor*, *propeller*, and *radio pulsar* phases, respectively. In case 1 we adopt the initial parameters as $M_d = 0.28 M_\odot$, $P_0 = 14 \text{ ms}$, $B_0 = 3 \times 10^{12} \text{ G}$, $M_0 = 10^{-5} M_\odot$,

and $R_d = 10^9$ cm. The NS is found to be spun-up to a period of 2.88 ms in 72 yr, when the disk becomes neutral¹. In the meanwhile, the magnetic field decreases dramatically to 4.2×10^8 G. This is a strongly spun-up pulsar, demonstrating a possible way for the formation of PSR J1903+0327. In case 2 we lower the initial disk mass M_d to $0.02M_\odot$, while keeping all the other parameters same as in case 1. This leads to a lower $\dot{M}_d(t_0)$, so that the spinning-up efficiency also becomes lower. The NS enters the *radio pulsar* phase with a relatively longer period of 11.26 ms and a higher magnetic field of 6.8×10^9 G. In case 3 we lower the initial disk radius R_d by a factor of 10 compared with in case 1. This gives a lower value of t_0 and a higher value of $\dot{M}_d(t_0)$, and the NS is spun-up to a period of 3.63 ms. In case 4 we choose a slower decay of the magnetic field than in case 1 by setting a higher value of $M_0 = 10^{-4} M_\odot$. It is shown that the period evolution remains almost unchanged, but the magnetic field decays to 4.2×10^9 G, about 10 times of that in case 1. A similar feature can be seen in case 5, if we increase the initial magnetic field to be $B = 1 \times 10^{13}$ G. Finally in case 6 M_d is further decreased to be $0.01 M_\odot$, now the NS is spun-up to a period of 11.82 ms during the first 39 yr and then spun-down during the *propeller* phase to a period of 11.92 ms during the subsequent 88 yr. After that time the disk becomes neutral and the *radio pulsar* phase begins. In the meanwhile the magnetic field decays to 6.6×10^{10} G. This is a mildly spun-up pulsar.

3. Population synthesis

Based on the analysis in the above section, now we try to estimate the statistical properties (e.g. the birth rate, total number, orbital period and eccentricity distributions) of RSR J1903+0327-like pulsars formed through the fallback disk accretion involved (FDAI) channel in the Galaxy, by using an evolutionary population synthesis (EPS) method based on the rapid binary stellar evolution (BSE) code developed by Hurley, Pols & Tout (2000) and Hurley, Tout & Pols (2002). This code incorporates the evolution of single stars with binary-star interactions, such as mass transfer, mass accretion, common-envelope (CE) evolution, SN kick, tidal friction, and angular momentum loss mechanisms (e.g. magnetic braking and gravitational radiation).

We assume all stars are born in binary systems. The initial mass function (IMF) of

¹The neutral time is given by $t_n \propto [R_d^{-3} \dot{M}_d(t_0)]^{1/2.75} \times R_d^{0.5}$ (Menou, Perna & Hernquist 2001). The value $t_n = 72$ yr used here is much less than that ($\sim 10^3 \sim 10^4$ yr) in Menou, Perna & Hernquist (2001). This is because (1) we calculated $\dot{M}_d(t_0)$ by normalizing it to the total disk mass, while Menou, Perna & Hernquist (2001) estimated it by assuming $\dot{M}_d(t_0) = M_d/t_0$, which gives a larger value of $\dot{M}_d(t_0)$ than ours; (2) the adopted value of $R_d = 10^9$ cm is larger than the value in Menou, Perna & Hernquist (2001).

Kroupa, Tout & Gilmore (1993) is adopted for the primary’s mass (M_1) distribution. For the secondary stars (M_2) we assume a uniform distribution of the mass ratio M_2/M_1 between 0 and 1. A uniform distribution of $\ln a$ is adopted for the binary separation a . The star formation rate parameter is taken to be $S = 7.6085 \text{ yr}^{-1}$, corresponding to a rate of $\sim 0.02 \text{ yr}^{-1}$ for core-collapse SNe in our Galaxy, assume all the stars with masses $> 8 M_\odot$ die through SN explosions.

It is known that a velocity kick is imparted to newborn NSs during core-collapse SNe, and a Maxwellian SN kick distribution with dispersion $V_\sigma \sim 200 - 400 \text{ km s}^{-1}$ is usually adopted (e.g. Lyne & Lorimer 1994; Hobbs et al. 2005). However, there is mounting evidence that some NSs may have received relatively small velocity kicks, of order less than 50 km s^{-1} (Pfahl, Rappaport & Podsiadlowski 2002; Pfahl et al. 2002; Podsiadlowski et al. 2004). A probable mechanism that causes small kicks is the electron capture (EC) SNe (Miyaji et al. 1980; Nomoto 1984). The explosion energies of such SNe are significantly lower than those of normal SNe (Dessart et al. 2006; Kitaura, Janka & Hillebrandt 2006; Jiang, Chen & Wang 2007), so that the nascent NSs receive a small kick velocity. It has been suggested that the initial masses of stars that explode via the EC SN mechanism may range in $\sim 6 - 12 M_\odot$ (Podsiadlowski et al. 2004; Kitaura, Janka & Hillebrandt 2006; Poelarends et al. 2008). Alternatively, core-collapse SN would be sub-energetic if the initial mass of its progenitor is greater than $\sim 18 M_\odot$. Simulations for a range of progenitor masses and different input physics suggest that the explosion energy decreases with increasing progenitor mass, whereas the binding energy of the star increases with increasing mass. This leads to a transition mass range of $\sim 18 - 25 M_\odot$ above which the SN explosion is so energy-deficient that fallback accretion would be important (Fryer 1999). For the aforementioned reasons, we assume a Maxwellian kick distribution with $V_\sigma = 20 \text{ km s}^{-1}$ for the progenitor stars with initial masses of $8 \leq M \leq 12 M_\odot$ and $M \geq 18 M_\odot$, otherwise we take $V_\sigma = 265 \text{ km s}^{-1}$.

The fallback disk mass M_d is poorly known, but various arguments suggest that $M_d \lesssim 0.1 M_\odot$ may not be unreasonable (Chevalier 1989; Lin, Woosley & Bodenheimer 1991). There should exist a relation between the disk mass and the properties of the progenitors. Unfortunately, there are big uncertainties in some key factors in the SN models, including the neutrino energy and transport, equation of state of gas at nuclear densities, convection, rotation, neutrino emission asymmetry, etc (Mönchmeyer 1991; Yamada & Sato 1995; Fryer 1999), each of which could strongly affect the SN energy and hence the mass of fallback. So we do not attempt to find an empirical formula to depict this kind of relation. Instead, we adopt simplified assumptions that sub-energetic SNe generally produce more massive fallback disks, i.e., the logarithm of the disk mass $\log(M_d/M_\odot)$ is distributed uniformly between -5 and -1 for stars of masses $8 \leq M \leq 12 M_\odot$ or $M \geq 18 M_\odot$, and between -6 to -2 in

other cases.

The initial magnetic fields are chosen so that $\log B$ is distributed normally with a mean of 12.4 and a standard deviation of 0.4. The magnetic field decay parameter M_0 is set to be $10^{-5} - 10^{-4} M_\odot$, and the disk radius $R_d \sim 10^7 - 10^8$ cm. The initial P_0 is chosen to be distributed uniformly between 10 and 100 ms. In this work an MPSR is defined as a radio pulsar with spin period less than 10 ms, and we *only* take the FDAI MPSRs (either in binaries or isolation) into account. In the case of BMPSRs, if the companion star fills its Roche lobe before the pulsar passes the death line, we take the elapsed time before this moment as the upper limit for the lifetime of the pulsar.

We divide the binary FDAI MPSRs into several categories according to their formation channels and the properties of the companion stars described as follows. The first case is that the pulsar is formed by core collapse of the primary star, and has an MS binary companion. Since the pulsar evolution can be strongly influenced by the stellar winds from the companion, which are intensive for massive stars, we only consider the MPSR-low-mass main sequence (MPSR-LMS) binaries, in which the companion star has a mass less than $1.5 M_\odot$. Alternatively the pulsar is formed by core collapse of the secondary, and already has an NS/BH binary companion. They are called NS/BH-MPSRs binaries. Finally, a single MPSR could form during either the first or the second SN that disrupts the binary.

To investigate the influence of the model parameters on the final results, we design nine models by changing the values of the CE parameter α_{CE} ², the field decay parameter M_0 , the lower and upper limit of the fallback disk mass $M_{d,low}$, $M_{d,upp}$ in case of sub-energetic SNe, and the disk radius R_d (listed in Table 2). The calculated numbers and formation rates of various types of binary and single MPSRs in our Galaxy are listed in Table 3, and briefly described as follows.

In model A (regarded as our control model) we adopt $\alpha_{CE} = 1.0$, $M_0 = 10^{-5} M_\odot$, $M_d = 10^{-5} - 0.1 M_\odot$, and $R_d = 10^8$ cm. It is predicted that in the Galaxy there are ~ 200 MPSR-LMS (PSR J1903+0327 may be one of them) and ten times more NS/BH-MPSR systems. Moreover, $\sim 10^5$ single MPSRs are produced. This number is comparable with estimated from observations (e.g. Lorimer 1995), suggesting that some fraction of the single MPSRs might form from the fallback disk accretion rather the traditional recycling channel in binary systems. Since their formation rate is 100 times lower than normal pulsars, it is

²The CE parameter α_{CE} is defined as the ratio of total binding energy of the envelope $E_{bind,i}$, and the difference between the initial and final orbital energies of the cores, i.e., $\alpha_{CE} = E_{bind,i}/(E_{orb,f} - E_{orb,i})$, where $E_{orb,f}$ and $E_{orb,i}$ are the final and initial binding energy of the cores, respectively (see Hurley, Tout & Pols 2002 for details).

not surprising that no MPSRs have been detected in young SN remnants.

In model B we increase the α_{CE} to 3.0, and find that higher α_{CE} produces more MPSR-LMS binaries. This is because binaries with a higher α_{CE} tend to survive during the CE evolution, especially for low-mass, close binaries which are more likely to be subject to coalescence during CE evolution. For the same reason, lower α_{CE} (model C) leads to fewer MPSRs (see Liu & Li 2006, for more detailed discussions). In model D we adopt a slower magnetic field decay, resulting in a shorter duration of the *accretor* phase, hence significantly reducing the numbers of all three types of MPSRs. In model E we choose a lower value of R_d than in model A. As mentioned in §2, this will cause longer final periods of the pulsars (compare the curves of P1 and P3 in Fig. 1), and produce fewer MPSRs. In model F the lower limit of the fallback disk masses is increased by a factor of 3, but the numbers of MPSRs change little, since MPSRs are produced only in the case of heavy disk accretion. This effect can also be illustrated in model G: when the upper limit of disk masses become half in model A, the numbers of MPSRs decrease by a factor of 10. To constrain the possible ranges of MPSR numbers, models H and I represent two extremely favorable and unfavorable cases for the production of MPSRs, respectively. It is found that the number of MPSR-LMS binaries may range from less than one to nearly one thousand in the Galaxy.

We plot the eccentricity vs. orbital period distributions of the MPSR-LMS binaries for models A, B, and C in Figure 2. The distributions of the other three models are similar as that of model A. It is seen that in both models A and C most MPSR-LMS binaries are in wide orbits ($P_{orb} > 10^3$ days) with eccentricities $e > 0.3$. For model B, the binary population seems to be distributed into two groups. One is similar as that in models A and C, the other stretches across large ranges of e ($\sim 0 - 1$) and P_{orb} ($\sim 1 - 10^3$ day) with a tendency of larger e accompanied with larger P_{orb} .

4. Discussion and Conclusions

In this paper we suggest a born-fast formation channel for MPSRs in the Galaxy. In this scenario, a newborn NS may experience spinning-up and field decay phase via accretion from a fossil disk, established as a result of fallback following a SN explosion. For appropriate choices of initial parameters, the NS could become a low field MPSR within $< 10^3$ yr. In particular, this scenario can naturally explain the properties (millisecond spin and eccentric orbit) of the recently discovered BMPSR PSR J1903+0327, without invoking stellar interactions within a GC or peculiar triple system.

Our population synthesis calculations also suggest a population of single FDAI MPSRs

in the Galaxy. The formation of single MPSRs has not been well understood, especially how the pulsar has lost its binary companion after mass transfer. The conventional explanation is that the pulsar was recycled in a binary located in a GC, and then ejected out due to stellar encounters, or the high-energy radiation from the pulsar has evaporated its companion (e.g. Ruderman, Shaham, & Tavani 1989). The latter explanation may also require a GC origin of the binary (King et al. 2005). Although the fraction of BMPSRs and LMXBs in GCs is much larger with respect to the fraction in the Galactic field, it seems unlikely that all single MPSRs have originated from GCs. The fallback disk accretion scenario might provide a possible way for the formation of not only single MSPs (Miller & Hamilton 2001) but also planetary systems around them (Lin, Woosley & Bodenheimer 1991).

Obviously the formation scenario proposed in this work is subject to many uncertainties, so the results in Table 3 should be regarded as the optimistic cases. One of the biggest issues is to determine how much fallback material would have enough angular momentum at the time of collapse to allow the formation of a disk. Our population synthesis calculations suggest that there may be tens to hundreds of MPSR-LMS binaries lurking in our Galaxy. For PSR J1903+0327, a relatively large amount of fallback disk mass ($\gtrsim 0.1 M_\odot$) is required to produce the very short period (< 3 ms), implying that the progenitor star may have experienced an unusual SN-fallback history. It is interesting to note that the large mass ($\sim 1.74 M_\odot$) of the pulsar may suggest a massive ($\gtrsim 18 M_\odot$) progenitor star (Zhang et al. 2008) and probably intensive fallback during the SN explosion. To form a centrifugally supported accretion disk around the NS, the specific angular momentum of the fallback matter should be larger than that needed for a circular orbit at the NS radius ($GM_{NS}R_{NS})^{1/2} \sim 1.4 \times 10^{16} \text{ cm}^2 \text{ s}^{-1}$. Modern stellar evolution calculations suggest, however, that the majority of massive stars lose angular momentum via magnetic processes very efficiently. Heger, Woosley & Spruit (2005) found that magnetic torques decrease the final rotation rate of the collapsing iron core by about a factor of 30 – 50 compared with the nonmagnetic counterparts. In their magnetic models there is a small amount of matter ($\sim 0.5 M_\odot$ for a $15 M_\odot$ star and $\sim 1.0 M_\odot$ for a $25 M_\odot$ star) that has specific angular momentum greater than $10^{16} \text{ cm}^2 \text{ s}^{-1}$ in the envelope of the pre-SN progenitor star. The realistic amount of fallback matter would be significantly less than these values because material with high specific angular momentum materials lies in the envelope of the progenitor star rather than in its inner core. Solutions of the problem of having enough rotation may involve tidal interactions in close binaries. If either star in the binary fills its Roche lobe, the spin-orbit coupling tidal torques would corotate the spins of two components with the orbital rotation, so that the progenitor star of the NS could be spun-up enough to allow the formation of the fallback disk. If this is true the realistic number of FDAI MPSRs should be less than our calculation, because not all the progenitors of the FDAI MPSRs have experienced the close binary evolution phases.

Other uncertainties include accretion from the SN fallback disk and magnetic field decay induced by accretion. The rate at which material falls back on the accretion disk depends on both the density profile of the star and on its angular velocity profile. Numerical calculations have been carried for a few mass and angular momentum configurations of massive stars (cf. Woosley & Bloom 2006, and references therein), which generally suggest highly super-Eddington accretion with a short time. Still lack is a comprehensive picture of the conditions for fallback disk formation after SN explosions of massive stars. Highly super-Eddington accretion in a short period of ~ 100 yr in a radiation-trapped regime plays a key role in the formation of FDAI MSPRs. This distinguishes our model from the standard type of accretion expected in the recycling scenario, which lasts for $\sim 10^8 - 10^{10}$ yr. In the standard pulsar recycling theory, the so called “spin-up line” in the $B - P$ diagram shows the minimum period to which the NS can be spun-up in Eddington-limited accretion. This line is defined by the equilibrium period assuming the spin-up proceeding at the Eddington accretion rate. Although the pulsars in our scenario have experienced the highly super-Eddington accretion, since the mass accretion rate declines rapidly, the equilibrium period is never attained, unlike in the case of accreting NS in LMXBs. Thus our model does not necessarily predict any MSPRs sitting above the spin-up line.

Nor is known how accretion-induced field decay occurs in NSs. One possible way is via ohmic dissipation of the accreting NSs crustal currents, due to the heating of the crust which in turn increases the resistance in the crust (Romani 1990; Geppert & Urpin 1994; Konar & Bhaattacharya 1997). Alternative scenarios consist of screening or burying the magnetic field with the accreted material (Bisnovatyi-Kogan & Komberg 1974; Taam & van den Heuvel 1986; Cumming et al. 2001), or outward moving vortices in the superfluid and superconducting core pushing magnetic fluxoids into the crust during pulsar’s spin-down (Srinivasan et al. 1990; Konar & Bhaattacharya 1999). In this work we require a fast field decay during the hyper-accretion phase (within $\lesssim 10^3 - 10^4$ yr). Screening or burying the magnetic field with the accreted material may be more appropriate for this scenario.

Finally we move to the uncertainties in the CE evolution. We have used a constant CE parameter α_{CE} to compute the orbital evolution during the spiral-in process, which is, however, very likely to change with the properties and evolutionary state of stars (Iben & Livio 1993; Taam & Sandquist 2000). From Fig. 2 we find that model B with $\alpha_{CE} = 3$ seems to be preferred for the formation of PSR J1903. This value is compatible with other estimates by, e.g., van den Heuvel (1994), Portegies Zwart & Yungelson (1998) and Kalogera (1999), but in contradiction with Taam (1996), Sandquist et al. (1998, 2000), and O’Shaughnessy (2008). Here the controversial issue is that there are no strict criteria for defining binding energy of stellar envelopes and there is no clear understanding whether sources other than gravitational energy may contribute to unbinding common envelopes (Han, Podsiadlowski & Eggleton

1995).

A distinct feature of the born-fast scenario is that the companion star of PSR J1903+0327 is predicted to be a “young” ($\lesssim 1 - 2 \times 10^9$ yr, i.e. within the characteristic age of the pulsar) MS star. Future optical, IR, and radio observations could present strong constraints on the nature of the optical counterpart, and thus verify or falsify the born-fast scenario.

We thank X. -P. Zheng, H. -L. Dai and Z. -Y. Zuo for helpful discussions, and J. R. Hurley for providing the BSE code. It is grateful to the anonymous referee for his/her insightful comments that help improving the original manuscript. This work was supported by the Natural Science Foundation of China under grant numbers 10573010 and 10221001. XWL thanks Huazhong Normal University for a postdoctoral fellowship.

REFERENCES

- Bhattacharya, D., & van den Heuvel, E. P. J. 1991, *Phys. Rep.*, 203, 1
- Bisnovatyi-Kogan, G. S. & Komberg, B.V. 1974, *SvA*, 18, 217
- Cannizzo, J. K., Lee, H. M., & Goodman, J. 1990, *ApJ*, 351, 38
- Champion, D. J., Ransom, S. M., Lazarus, P., Camilo, F., Bassa, C. et al. 2008, *Science*, 320, 1309
- Chatterjee, P., Hernquist, L., & Narayan, R. 2000, *ApJ*, 534, 373
- Chevalier, R. A. 1989, *ApJ*, 346, 847
- Colgate, S. A. 1971, *ApJ*, 163, 221
- Cumming, A., Zweibel, E., & Bildsten, L. 2001, *ApJ*, 557, 958
- Dessart, L., Burrows, A., Ott, C. D., Livne, E., Yoon, S. -C., & Langer, N. 2006, *ApJ*, 644, 1063
- Francischelli, G. J., & Wijers, R. A. M. J. 2002, *ApJ*,
- Freire, P. C. C., Ransom, S. M., & Gupta, Y. 2007, in the proceedings of ”40 Years of Pulsars: Millisecond Pulsars, Magnetars, and More”, August 12-17, 2007, McGill University, Montreal, Canada, [arXiv:0711.1883]
- Fryer, C. L. 1999, *ApJ*, 522, 413

- Geppert, U. & Urpin, V. 1994, MNRAS, 271, 490
- Han, Z. W., Podsiadlowski, P., & Eggleton, P. P. 1995, MNRAS, 272, 800
- Heger, A., Woosley, S. E., & Spruit, H. C. 2005, ApJ, 626, 350
- Hobbs, G., Lorimer, D. R., Lyne, A. G., & Kramer, M. 2005, MNRAS, 360, 974
- Hurley, J. R., Pols, O. R., & Tout, C. A. 2000, MNRAS, 315, 543
- Hurley, J. R., Tout, C. A., & Pols, O. R. 2002, MNRAS, 329, 897
- Iben, I. Jr. & Livio, M. 1993, PASP, 105, 1373
- Illarinov, A. F., & Sunyaev, R. A. 1975, A&A, 39, 185
- Jiang, B., Chen, Y., & Wang, Q. D. 2007, ApJ, 670, 1142
- Jiang, Z. -B., Li, X. -D. 2005, ChJAA, 5, 487
- Kalogera, V. 1999, ApJ, 521, 723
- King, A. R., Beer, M. E., Rolfe, D. J., Schenker, K., & Skipp, J. M. 2005, MNRAS, 358, 1501
- Kitaura, F. S., Janka, S. T., & Hillebrandt, W. 2006, A&A, 450, 345
- Konar, S. & Bhaattacharya, D., 1997, MNRAS, 284, 311
- Konar, S. & Bhattacharya, D. 1999, MNRAS, 308, 795
- Kroupa, P., Tout, C. A., & Gilmore, G. 1993, MNRAS, 262, 545
- Li, X. -D. 2002, ApJ, 579, L37
- Lin, D. N. C., Woosley, S. E., & Bodenheimer, P. H. 1991, Nature, 353, 827
- Liu, X. -W. & Li, X. -D. 2006, A&A, 449, 135
- Lorimer, D. R. 1995, MNRAS, 274, 300
- Lyne, A. G., & Lorimer, D. R. 1994, Nature, 369, 127
- Menou, K., Perna, R., & Hernquist 2001, ApJ, 559, 1032
- Miller, M. C. & Hamilton, D. P. 2001, ApJ, 550, 863

- Miyaji, S., Nomoto, K., Yokoi, K., & Sugimoto, D. 1980, PASJ, 32, 303
- Mönchmeyer, R. 1991, in *Supernovae*, The Tenth Santa Cruz Workshop in Astronomy and Astrophysics, ed. Woosley, S. E., 358
- Nomoto, K. 1984, ApJ, 277, 791
- O’Shaughnessy, R., Kim, C., Kalogera, V., & Belczynski, K. 2008, ApJ, 672, 479
- Pfahl, E., Rappaport, S., & Podsiadlowski, P. 2002, ApJ, 571, 37
- Pfahl, E., Rappaport, S., Podsiadlowski, P., & Spruit, H. 2002, ApJ, 574, 364
- Podsiadlowski, P., Langer, N., Poelarends, A. J. T., Rappaport, S., Heger, A., Pfahl, E. 2004, ApJ, 612, 1044
- Poelarends, A. J. T., Herwig, F., Langer, N., & Heger, A. 2008, ApJ, 675, 614
- Portegies Zwart, S. F. & Yungelson, L. R. 1998, A&A, 332, 173
- Romani, R. W. 1990, Nature, 347, 741
- Ruderman, M., Shaham, J., & Tavani, M. 1989, ApJ, 336, 507
- Ruderman, M., & Sutherland P. G. 1975, ApJ, 196, 51
- Sandquist, E. L., Taam, R. E., & Burkert, A. 2000, ApJ, 533, 984
- Sandquist, E. L., Taam, R. E., Chen, X., Bodenheimer, P., & Burkert, A. 1998, ApJ500, 909
- Shibazaki, N., Murakami, T., Shaham, J., Nomoto, K. 1989, Nature, 342, 656
- Srinivasan, G., Bhattacharya, D., Muslimov, A. G., & Tsygan, A. J. 1990, Current Science, 59, 3
- Taam, R. E. 1996. In *Compact Stars in Binaries*, eds. J. van Paradijs, E. P. J. van den Heuvel, E. Kuulkers (Dordrecht: Kluwer), p. 3
- Taam, R. E., & Sandquist, E. L. 2000, ARA&A, 38, 113
- Taam, R. E., & van den Heuvel, E. P. J. 1986, ApJ, 305, 235
- bibitem[Tauris & van den Heuvel(2006)]tauris06 Tauris, T. M., & van den Heuvel, E. P. J. 2006 in *Compact stellar X-ray sources*. Edited by Walter Lewin & Michiel van der Klis. Cambridge Astrophysics Series, No. 39. Cambridge, UK: Cambridge University Press, p.623

- van den Heuvel, E. P. J. 1994, A&A, 291, L39
- van den Heuvel, E. P. J. 2008, Science, 320, 1298
- Woosley, S. E. & Bloom, J. S. 2006, ARA&A, 44, 507
- Yamada, S., & Sato, K. 1995, ApJ, 434, 268
- Zhang, W., Woosley, S. E., & Heger, A. 2008, ApJ, 679, 639




Fig. 1.— Period (bottom-left axes) and magnetic field (top-right axes) evolution of an NS with initial parameters list in Table 1. The curves labeled P1 - P6 and B1 - B6 represent the evolutions of period and magnetic field, respectively. Numbers are suffixed to indicate the corresponding models. In the $P - t$ plot, the thick solid lines, solid lines, and dotted lines represent the *accretor*, the *propeller*, and the *radio pulsar* phases, respectively. The magnetic field evolution is plotted until the *radio pulsar* phase begins.




Fig. 2.— Orbital period and eccentricity probability distribution of the MPSR-LMS binaries for model A, B, and C, from left to right, respectively. The distributions for the other six models (D-G, H, and I) are very similar to those for models A, B, and C.

Table 1: Initial model parameters for the NS-disk systems.

Model No.	M_d ($M_\odot$)	$R_{d,10}$	P_0 (ms)	B_0 (10^{12} G)	M_0 ($10^{-5} M_\odot$)
1	0.28	0.1	14	3	1
2	0.02	0.1	14	3	1
3	0.28	0.01	14	3	1
4	0.28	0.1	14	3	10
5	0.28	0.1	14	10	1
6	0.01	0.01	14	10	1

Table 2: Model parameters for binary population synthesis.

Model	α_{CE}	M_0 ($10^{-5} M_\odot$)	$R_{d,8}$	$M_{d,low}$ ($10^{-5} M_\odot$)	$M_{d,upp}$ ($0.1 M_\odot$)
A	1	1	1	1	1
B	3	1	1	1	1
C	0.5	1	1	1	1
D	1	10	1	1	1
E	1	1	0.1	1	1
F	1	1	1	3	1
G	1	1	1	1	0.5
H	3	1	1	3	1
I	0.5	10	0.1	1	0.5

Table 3: Predicted numbers and formation rates in our Galaxy of various types of FDAI MPSR systems.

Model	item	MPSR-LMS	NS/BH-MPSR	Single MPSR
A	number	203	3.3×10^3	1.5×10^5
	rate	6.0×10^{-7}	3.1×10^{-6}	3.1×10^{-4}
B	number	751	3.7×10^3	1.6×10^5
	rate	1.3×10^{-6}	4.0×10^{-6}	3.2×10^{-4}
C	number	192	3.1×10^3	1.5×10^5
	rate	6.5×10^{-7}	3.0×10^{-6}	3.1×10^{-4}
D	number	8	1.4×10^2	1.2×10^3
	rate	6.2×10^{-7}	3.3×10^{-6}	2.7×10^{-4}
E	number	83	1.3×10^3	4.9×10^4
	rate	3.9×10^{-7}	2.0×10^{-6}	1.9×10^{-4}
F	number	232	3.7×10^3	1.7×10^5
	rate	6.9×10^{-7}	3.5×10^{-6}	3.4×10^{-4}
G	number	28	4.5×10^2	1.2×10^4
	rate	3.5×10^{-7}	1.7×10^{-6}	2.0×10^{-4}
H	number	856	4.2×10^3	1.8×10^5
	rate	1.5×10^{-6}	4.5×10^{-6}	3.6×10^{-4}
I	number	0.3	7	38
	rate	9.9×10^{-8}	4.6×10^{-7}	3.9×10^{-5}